

LEFT REGULAR REPRESENTATION OF $sl_q(3)$:**REDUCTION AND INTERTWINERS****Ludwik Dąbrowski and Preeti Parashar**

SISSA, Strada Costiera 11, 33014 Trieste, Italy

Abstract

Reduction of the left regular representation of quantum algebra $sl_q(3)$ is studied and q -difference intertwining operators are constructed. The irreducible representations correspond to the spaces of local sections of certain line bundles over the q -flag manifold.

1. Introduction.

Recently, following the canonical procedure for $q = 1$ [1], the left regular representation, its reduction to infinite family of (reducible and irreducible) representations and the q -difference intertwining operators were studied on the examples of Lorentz quantum algebra [2], of $sl_q(2)$ and its contraction to $e_q(2)$ [3]. In this paper we investigate the case of $sl_q(3)$, a quantization of the simple Lie algebra of rank two. This Hopf algebra, besides being computationally more involved, presents some new features which are interesting enough in order to generalize to the case of $sl_q(n)$ and next to q -deformations of all semisimple Lie algebras. This will help to understand better the important relation between representation theory and geometry of quantum groups.

Denote by G_q the quantum group $\text{Fun}_q(G)$ and by g_q its dual $U_q(g)$, a quantization of the Lie algebra g of G . Recall that the left regular representation $L : g_q \times G_q \rightarrow G_q$, $(a, \varphi) \rightarrow \mathcal{L}(a)\varphi$, and the right regular representation $R : g_q \times G_q \rightarrow G_q$, $(a, \varphi) \rightarrow \mathcal{R}(a)\varphi$, of g_q are defined on G_q by, respectively

$$(\mathcal{L}(a)\varphi)(b) = \varphi(S(a)b) \tag{1.1}$$

$$(\mathcal{R}(a)\varphi)(b) = \varphi(ba), \tag{1.2}$$

where $a, b \in g_q$, $\varphi \in G_q$ and S is the antipode.

We shall reduce the left regular representation on the eigenspaces of the right regular representation of the Borel generators. Namely, we shall impose the condition that the Cartan generators act as a multiplication by fixed numbers (to be specified further) and that the other Borel generators vanish. Then some of the remaining generators of g yield (via the right regular representation) interesting intertwining q -difference operators, the kernels of which provide a further reduction. This method yields a natural q -deformation of several equations of mathematical physics, which possess a symmetry group and which can be viewed as intertwiner between some representations of the Lie algebra of the symmetry. It may be helpful also in the inverse problem of finding the symmetry behind a given q -difference equation.

2. Preliminaries on $sl_q(3)$ and $SL_q(3)$.

The generators of $SL_q(3)$ (quantization of the algebra of complex functions on $SL_q(3)$) are the unit 1 and t_{ij} , with $i, j = 1, 2, 3$; arranged as a 3×3 matrix T , satisfying the commutation relations

$$R T_1 T_2 = T_2 T_1 R \quad (T_1 = T \otimes I, T_2 = I \otimes T), \quad (2.1)$$

where

$$q^{\frac{1}{3}} R = q \sum_i e_{ii} \otimes e_{ii} + \sum_{i \neq j} e_{ii} \otimes e_{jj} + \lambda \sum_{i > j} e_{i,j} \otimes e_{j,i}, \quad (2.2)$$

with $\lambda = q - q^{-1}$. More explicitly, we have:

$$t_{11}t_{12} = qt_{12}t_{11}, \quad t_{11}t_{13} = qt_{13}t_{11}, \quad t_{11}t_{21} = qt_{21}t_{11}, \quad (2.3)$$

$$t_{11}t_{31} = qt_{31}t_{11}, \quad [t_{11}, t_{23}] = \lambda t_{13}t_{21}, \quad [t_{11}, t_{32}] = \lambda t_{12}t_{32},$$

$$t_{22}t_{12} = q^{-1}t_{12}t_{22}, \quad t_{22}t_{13} = t_{13}t_{22}, \quad t_{22}t_{23} = qt_{23}t_{22}, \quad (2.4)$$

$$t_{22}t_{21} = q^{-1}t_{21}t_{22}, \quad t_{22}t_{31} = t_{31}t_{22}, \quad t_{22}t_{32} = qt_{32}t_{22},$$

$$[t_{33}, t_{12}] = -\lambda t_{13}t_{32}, \quad t_{33}t_{13} = q^{-1}t_{13}t_{33}, \quad t_{33}t_{23} = q^{-1}t_{23}t_{33}, \quad (2.5)$$

$$[t_{33}, t_{21}] = -\lambda t_{23}t_{31}, \quad t_{33}t_{31} = q^{-1}t_{31}t_{33}, \quad t_{33}t_{32} = q^{-1}t_{32}t_{33},$$

$$[t_{11}, t_{22}] = \lambda t_{12}t_{21}, \quad [t_{11}, t_{33}] = \lambda t_{13}t_{31}, \quad [t_{22}, t_{33}] = \lambda t_{23}t_{32}, \quad (2.6)$$

$$t_{12}t_{13} = qt_{13}t_{12}, \quad t_{12}t_{21} = t_{12}t_{21}, \quad [t_{12}, t_{23}] = \lambda t_{13}t_{22}, \quad t_{12}t_{31} = t_{31}t_{12}, \quad t_{12}t_{32} = qt_{32}t_{12}, \quad (2.7)$$

$$t_{13}t_{21} = t_{21}t_{13}, \quad t_{13}t_{23} = qt_{23}t_{13}, \quad t_{13}t_{31} = t_{31}t_{13}, \quad t_{13}t_{32} = t_{32}t_{13}, \quad (2.8)$$

$$t_{21}t_{23} = qt_{23}t_{21}, \quad t_{21}t_{31} = qt_{31}t_{21}, \quad [t_{21}, t_{32}] = \lambda t_{31}t_{22}, \quad (2.9)$$

$$t_{23}t_{31} = t_{31}t_{23}, \quad t_{23}t_{32} = t_{32}t_{23}, \quad (2.10)$$

$$t_{31}t_{32} = qt_{32}t_{31}. \quad (2.11)$$

An additional relation is

$$Det_q = t_{11}(t_{22}t_{33} - qt_{23}t_{32}) - qt_{21}(t_{12}t_{33} - qt_{13}t_{32}) + q^2t_{31}(t_{12}t_{23} - qt_{13}t_{22}) = 1 \quad (2.12)$$

which can be written also in equivalent form as

$$Det_q = t_{11}(t_{22}t_{33} - qt_{23}t_{32}) - qt_{12}(t_{21}t_{33} - qt_{23}t_{31}) + q^2t_{13}(t_{21}t_{32} - qt_{22}t_{31}) = 1 \quad (2.13)$$

Considered as a Hopf algebra, $SL_q(3)$ has the following comultiplication Δ , counit ε , and antipode S given on the generators by:

$$\Delta t_{ij} = t_{ik} \otimes t_{kj} \quad (2.14)$$

$$\varepsilon t_{ij} = \delta_{ij} \quad (2.15)$$

$$St_{ij} = (T^{-1})_{ij}. \quad (2.16)$$

For $\bar{q} = q$, with the $*$ -conjugation (complex anti-linear algebra anti-involution and co-algebra involution) $t_{ij}^* = St_{ji}$, $SL_q(3)$ becomes a Hopf $*$ -algebra denoted $SU_q(3)$.

As generators of the algebra $sl_q(3)$, dual to $SL_q(3)$, we shall use the unit 1 and the functionals which have been introduced in [4]: l_{ij}^+ with $i < j$; l_{ij}^- with $i > j$; l_{ii}^+ and l_{ii}^- , where $i, j = 1, 2, 3$. When arranged in upper- and lower-triangular matrices $L^\pm$, respectively, they are defined by the duality conditions

$$(L^\pm, T_1 \dots T_m) = R_1^\pm \dots R_m^\pm, \quad \text{for } m = 1, 2, \dots; \quad (2.17)$$

where for $1 \leq \ell \leq m$, $T_\ell^\pm$ act in the ℓ^{th} factor; $R_\ell^\pm$ act in the $(0, \ell)^{th}$ factor of $(\mathcal{C}^3)^{\otimes(m+1)}$; $R^+ = PRP$, $R^- = R^{-1}$ and $P \in \text{Mat}(\mathcal{C}^3 \otimes \mathcal{C}^3)$ is the permutation matrix.

The commutation relations are

$$R^+ L_1^\pm L_2^\pm = L_2^\pm L_1^\pm R^+, \quad R^+ L_1^+ L_2^- = L_2^- L_1^+ R^+. \quad (2.18)$$

Additional relations are:

$$l_{ii}^+ l_{ii}^- = 1, \quad i = 1, 2, 3; \quad (2.19)$$

$$l_{11}^+ l_{22}^+ l_{33}^+ = l_{11}^- l_{22}^- l_{33}^- = 1. \quad (2.20)$$

The Hopf algebra structure is then given by

$$\Delta l_{ij}^\pm = l_{ik}^\pm \otimes l_{kj}^\pm \quad (2.21)$$

$$\varepsilon l_{ij}^\pm = \delta_{ij} \quad (2.22)$$

and the antipode which is defined as for T with the replacement of q by q^{-1} . The $l_{ij}^\pm$ can be expressed in terms of more popular generators q^{H_i} , $X_i^\pm$, $i = 1, 2$ via:

$$\begin{aligned} q^{H_1} &= l_{11}^+ (l_{22}^+)^{-1}, \quad q^{H_2} = l_{11}^+ (l_{22}^+)^2, \\ X_1^+ &= \lambda^{-1} q^{-1/6} l_{12}^+ (l_{11}^+)^{-1/2} (l_{22}^+)^{-1/2}, \quad X_2^+ = \lambda^{-1} q^{-1/6} l_{23}^+ (l_{11}^+)^{1/2}, \\ X_1^- &= -\lambda^{-1} q^{1/6} l_{21}^- (l_{11}^+)^{1/2} (l_{22}^+)^{1/2}, \quad X_2^- = -\lambda^{-1} q^{1/6} l_{32}^- (l_{11}^+)^{-1/2}. \end{aligned} \quad (2.23)$$

They satisfy the following commutation relations:

$$\begin{aligned}
[H_i, H_j] &= 0, \quad [H_i, X_j^\pm] = \pm(\alpha_i, \alpha_j) X_j^\pm, \\
[X_i^+, X_j^-] &= \delta_{ij} [H_i], \quad i, j = 1, 2; \\
\sum_{k=0}^m (-1)^k \binom{m}{k}_q q^{k(k-m)(\alpha_i, \alpha_i)/2} (X_i^\pm)^k X_j^\pm (X_i^\pm)^{m-k} &= 0 \text{ for } i \neq j,
\end{aligned} \tag{2.24}$$

where $[H_i] = \frac{(q^{H_i} - q^{-H_i})}{(q - q^{-1})}$, $m = 1 - A_{ij}$ and $\binom{m}{k}_q = \frac{(q^m - 1)(q^{m-1} - 1) \dots (q^{m-k+1} - 1)}{(q^k - 1)(q^{k-1} - 1) \dots (q - 1)}$.

The Hopf algebra structure is then given by

$$\Delta(H_i) = H_i \otimes 1 + 1 \otimes H_i, \quad \Delta(X_i^\pm) = X_i^\pm \otimes q^{-H_i/2} + q^{H_i/2} \otimes X_i^\pm \tag{2.25}$$

$$\varepsilon(X_i^\pm) = 0, \quad \varepsilon(H_i) = 0 \tag{2.26}$$

$$S(X_i^\pm) = q^{-\rho} X_i^\pm q^\rho, \quad S(H_i) = -H_i \tag{2.27}$$

with $\rho = \sum H_\alpha/2$, where α belongs to the set of positive roots.

3. The left regular representation of $sl_q(3)$ on $SL_q(3)$.

It can be easily seen from the definition (1.1) that the left regular representation of $l_{ij}^\pm$ amounts to a multiplication of the generator matrix T from the left by, respectively, the numerical matrices :

$$\begin{aligned}
(Sl_{11}^+, T) &= q^{\frac{1}{3}} \begin{pmatrix} \frac{1}{q} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, (Sl_{12}^+, T) = -q^{\frac{1}{3}} \lambda \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, (Sl_{13}^+, T) = -q^{\frac{1}{3}} \lambda \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \\
(Sl_{21}^-, T) &= q^{-\frac{1}{3}} \lambda \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, (Sl_{22}^+, T) = q^{\frac{1}{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{q} & 0 \\ 0 & 0 & 1 \end{pmatrix}, (Sl_{23}^+, T) = -q^{\frac{1}{3}} \lambda \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \\
(Sl_{31}^-, T) &= q^{-\frac{1}{3}} \lambda \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, (Sl_{32}^-, T) = q^{-\frac{1}{3}} \lambda \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, (Sl_{33}^+, T) = q^{\frac{1}{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \frac{1}{q} \end{pmatrix}.
\end{aligned} \tag{3.1}$$

Similarly, the right regular representation amounts to a multiplication of T from the right by, respectively,

$$\begin{aligned}
(T, l_{11}^+) &= q^{-\frac{1}{3}} \begin{pmatrix} q & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, (T, l_{12}^+) = q^{-\frac{1}{3}} \lambda \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, (T, l_{13}^+) = q^{-\frac{1}{3}} \lambda \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \\
(T, l_{21}^-) &= -q^{\frac{1}{3}} \lambda \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, (T, l_{22}^+) = q^{-\frac{1}{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & q & 0 \\ 0 & 0 & 1 \end{pmatrix}, (T, l_{23}^+) = q^{-\frac{1}{3}} \lambda \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \\
(T, l_{31}^-) &= -q^{\frac{1}{3}} \lambda \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, (T, l_{32}^-) = -q^{\frac{1}{3}} \lambda \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, (T, l_{33}^+) = q^{-\frac{1}{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & q \end{pmatrix}.
\end{aligned} \tag{3.2}$$

As an (overcomplete) basis of $SL_q(3)$ we can take the following ordered monomials:

$$t^{\bar{n}} =: t_{11}^{n_{11}} t_{12}^{n_{12}} \dots t_{33}^{n_{33}}, \tag{3.3}$$

where $\bar{n} =: (n_{11}, n_{12}, \dots, n_{33})$ with $n_{ij} \in \mathbb{Z}_+$ and we use (unless otherwise stated) the usual ordering, i.e. first according to the row index and, then, the column index.

In the sequel we shall need an *independent* basis for $SL_q(3)$ which we believe must be known, but we have not been able to find it in the literature for $SL_q(n)$ with $n \geq 3$. In order to obtain such an independent basis we have to restrict the elements $t^{\bar{n}}$ by the $\det_q = 1$ condition (2.2). We proceed as follows.

Assume for a moment another ordering according to which the maximal power $(t_{11}t_{22}t_{33})^p$, where $p = \min\{n_{11}, n_{22}, n_{33}\}$, of product of diagonal generators t_{11} , t_{22} and t_{33} come first, then the remaining powers of (at most two of) t_{ii} and finally the other generators t_{ij} , $i \neq j$, in the usual order. (This is a convenient choice, inessential for the outcome). Using (2.2), $t_{11}t_{22}t_{33}$ can be expressed as a polynomial in t_{ij} of degree at most one in any of the generators t_{11} , t_{22} and t_{33} . Unfortunately, by reordering we may raise the degree in the diagonal generators. However, by analysing the commutation relations (2.3)-(2.11) we see that some of them (of the exchange type) are innoquous. Those which are of the commutator type and between the diagonal generators produce only the non-diagonal generators t_{ij} and lower the degree of two diagonal generators. Next, those of the commutator type between the non-diagonal generators produce at most one diagonal generator (namely t_{22}). Altogether, there is a net lowering of the complexive degree of the product $t_{11}t_{22}t_{33}$. By iteration, this allows to lower consistently this degree up to reaching

the situation when one of the diagonal generators is not present. Finally, by excluding the overlapping cases we arrive at the following result:

The independent basis in $sl_q(3)$ consist of three sectors composed by the elements $t^{\bar{n}}$ with one of the following restrictions:

$$\begin{aligned} (I) \quad & n_{11} = 0 \\ (II) \quad & n_{11} \geq 1, n_{22} = 0 \\ (III) \quad & n_{11} \geq 1, n_{22} \geq 1, n_{33} = 0 . \end{aligned} \tag{3.4}$$

Now, by iterating the twisted derivation rules for $sl_q(3)$

$$\mathcal{L}(l_{ij}^{\pm})\varphi\psi = \sum_k \mathcal{L}(l_{kj}^{\pm})\varphi \cdot \mathcal{L}(l_{ik}^{\pm})\psi , \tag{3.5}$$

$$\mathcal{R}(l_{ij}^{\pm})\varphi\psi = \sum_k \mathcal{R}(l_{ik}^{\pm})\varphi \cdot \mathcal{R}(l_{kj}^{\pm})\psi , \tag{3.6}$$

we obtain the action of the representation $\mathcal{L}$ on $t^{\bar{n}}$ which we give in the Appendix.

4. Reduction and intertwiners.

We impose the conditions of the (infinitesimal) right covariance on the independent basis for $SL_q(3)$. First, we consider only those functions ψ (i.e. the multilabels $\bar{n}$) on which

$$\mathcal{R}(l_{ij}^+) \psi = 0 , \quad i < j . \tag{4.1}$$

By these conditions some of the indices n_{ij} have to vanish, but for some other there is a possibility of compensation between various pieces, c.f. (A 10-12). This requires solutions to be of the form of infinite series. Instead, we shall assume invertibility of some (special combinations of) generators t_{ij} and may recover the series by formal expansion of the inverses in q . By a deeper look, we see that the solutions of (4.1) have to be built from generators t_{i3} in the last column and from minors m_{i1} of the first column (m_{ij} is the minor of t_{ij} , i.e. the q -determinant of T with i^{th} line and j^{th} column removed). Due to (2.12) one of them is dependent on the rest and we choose to delete m_{31} . Next, we impose that on ψ

$$\mathcal{R}(l_{ii}^+) \psi = \exp r_i \psi , \tag{4.2}$$

for $i = 1, 2$ (the case $i = 3$ is not independent due to (2.20)), where the labelling numbers r_1 and of r_2 will be specified further. We note the multiplicative property of the eigenvalue of $\mathcal{R}(l_{ii}^+)$ with respect to the product of eigenvectors and note that consequently the quotients $t_{i3}t_{j3}^{-1}$ and $m_{i1}m_{j1}^{-1}$ have eigenvalue 1. Among them, we choose to work with the following three independent combinations:

$$w_1 = t_{13}t_{33}^{-1}, \quad w_2 = t_{23}t_{33}^{-1}, \quad w_3 = m_{21}m_{11}^{-1}, \quad (4.3)$$

where $m_{21} = t_{12}t_{33} - qt_{13}t_{32}$ and $m_{11} = t_{22}t_{33} - qt_{23}t_{32}$.

Instead, in order to solve (4.2) , we employ the combination $m_{11}^{j_1} t_{33}^{j_2}$, with j_1, j_2 to be determined in terms of r_1, r_2 . Indeed, we get

$$r_1 = -(j_2 + 2j_1)h/3, \quad r_2 = -(j_1 + j_2)h/3,$$

where $q = \exp h$. Thus, since $j_1, j_2 \in \mathbb{Z}$, the admissible values of r_1 and of r_2 have to be quantized in units of $h/3$. Finally, the basis in the space $\mathcal{T}_{\bar{j}}$, $\bar{j} =: (j_1, j_2)$, of common solutions of (4.1) and (4.2) consist of the ordered monomials:

$$\phi_{\bar{j}}^{\bar{n}} =: w_1^{n_1} w_2^{n_2} w_3^{n_3} m_{11}^{j_1} t_{33}^{j_2}, \quad (4.4)$$

where $n_1, n_2, n_3 \in \mathbb{Z}_+$. In this way we obtain an infinite family of reduced representation spaces $\mathcal{T}_{\bar{j}}$, indexed by the integers j_1, j_2 .

It is remarkable, that the three independent variables w_i form a closed algebra:

$$w_1 w_2 = q w_2 w_1, \quad w_1 w_3 = q^{-1} w_3 w_1, \quad w_2 w_3 = q w_3 w_2 + \lambda w_1. \quad (4.5)$$

We give also some relations useful for the following computations:

$$w_3^n w_2 = q^{-n} w_2 w_3^n + \lambda q^{-1} [n] w_1 w_3^{n-1}, \quad w_3 w_2^n = q^{-n} w_2^n w_3 + \lambda q^{-n} [n] w_1 w_2^{n-1}. \quad (4.6)$$

Moreover, the two ‘spectators’ t_{33} and m_{11} commute:

$$t_{33} m_{11} = m_{11} t_{33} \quad (4.7)$$

and have the following commutation relations with the variables w_i :

$$w_1 m_{11} = q m_{11} w_1, \quad w_2 m_{11} = m_{11} w_2, \quad w_3 m_{11} = q m_{11} w_3, \quad (4.8)$$

$$w_1 t_{33} = q t_{33} w_1, \quad w_2 t_{33} = q t_{33} w_2, \quad w_3 t_{33} = t_{33} w_3. \quad (4.9)$$

We note that our w_i are also the relevant variables as far as the global covariance is concerned. This can be seen from the Gauss decomposition of T :

$$\begin{pmatrix} 1 & w_3 & w_1 \\ 0 & 1 & w_2 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} t_{11} - w_3 t_{33}^{-1} m_{12} - w_1 t_{31} & 0 & 0 \\ 0 & t_{33}^{-1} m_{11} & 0 \\ 0 & 0 & t_{33} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ m_{11}^{-1} m_{12} & 1 & 0 \\ t_{33}^{-1} t_{31} & t_{33}^{-1} t_{32} & 1 \end{pmatrix}. \quad (4.10)$$

By a lengthy but straightforward computation, iterating the twisted derivation rules, we obtain the explicit formulas for the representation in $\mathcal{T}_{\bar{j}}$:

$$\mathcal{L}(l_{12}^+) \phi_{\bar{j}}^{\bar{n}} = -\lambda q^{1-n_2-n_3+(2j_1+j_2)/3} [n_2] \phi_j^{n_1, n_2-1, n_3} + \lambda q^{1-n_2+(j_2-j_1)/3} [n_3-j_1] \phi_j^{n_1, n_2, n_3+1} \quad (4.11)$$

$$\mathcal{L}(l_{23}^+) \phi_{\bar{j}}^{\bar{n}} = \lambda q^{1-(2j_2+j_1)/3} [n_1+n_2-n_3-j_2] \phi_j^{n_1, n_2+1, n_3} + \lambda q^{n_1-(2j_2+j_1)/3} [n_3] \phi_j^{n_1+1, n_2, n_3-1} \quad (4.12)$$

$$\begin{aligned} \mathcal{L}(l_{13}^+) \phi_{\bar{j}}^{\bar{n}} &= \lambda q^{1-n_2+(4j_2-j_1)/3} [n_1-n_2-n_3-j_1+j_2] \phi_j^{n_1+1, n_2, n_3} \\ &+ \lambda q^{2-n_1-n_2+(j_2-j_1)/3} [j_1-n_3] \phi_j^{n_1, n_2+1, n_3+1} \end{aligned} \quad (4.13)$$

$$\mathcal{L}(l_{11}^+) \phi_{\bar{j}}^{\bar{n}} = q^{-n_1-n_3+(2j_1+j_2)/3} \phi_{\bar{j}}^{\bar{n}} \quad (4.14)$$

$$\mathcal{L}(l_{22}^+) \phi_{\bar{j}}^{\bar{n}} = q^{-n_2+n_3+(j_2-j_1)/3} \phi_{\bar{j}}^{\bar{n}} \quad (4.15)$$

$$\mathcal{L}(l_{33}^+) \phi_{\bar{j}}^{\bar{n}} = q^{n_1+n_2-(j_1+2j_2)/3} \phi_{\bar{j}}^{\bar{n}} \quad (4.16)$$

$$\mathcal{L}(l_{21}^-) \phi_{\bar{j}}^{\bar{n}} = \lambda q^{n_1-1+(j_1-j_2)/3} [n_3] \phi_j^{n_1, n_2, n_3-1} + \lambda q^{n_2-n_3+(j_1-j_2)/3} [n_1] \phi_j^{n_1-1, n_2+1, n_3} \quad (4.17)$$

$$\mathcal{L}(l_{32}^-) \phi_{\bar{j}}^{\bar{n}} = \lambda q^{-1+(j_1+2j_2)/3} [n_2] \phi_j^{n_1, n_2-1, n_3} \quad (4.18)$$

$$\mathcal{L}(l_{31}^-) \phi_{\bar{j}}^{\bar{n}} = \lambda q^{n_2-1+(j_1+2j_2)/3} [n_1] \phi_j^{n_1-1, n_2, n_3} . \quad (4.19)$$

So far we have restricted the left regular representation to (infinite dimensional) subrepresentations $\mathcal{T}_{\bar{j}}$. The restricted functions depend effectively only on the variables w_i since for each fixed $\bar{j}$ the factor $m_{11}^{j_1} t_{33}^{j_2}$ is fixed. In order to pursue the reduction till the end and get the functions depending *manifestly* only on w_i we associate with each $\phi \in T_{\bar{j}}$ a function $\hat{\phi}$, by the formula

$$\hat{\phi} = \phi m_{11}^{-j_1} t_{33}^{-j_2} . \quad (4.20)$$

We note that we have actually a freedom to work with variables proportional to w_i ,

$$z_i = C_i w_i, \quad (4.21)$$

where C_i are constants depending on q and $\bar{j}$. (This leads to an equivalent representation). In order to simplify the formulae we choose

$$C_1 = C_2 = q^{1-(j_1+2j_2)/3}, \quad C_3 = q^{1+(j_2-j_1)/3}. \quad (4.22)$$

The representations (4.11)- (4.19) acting in $\mathcal{T}_{\bar{j}}$, induce the following transformation rules on the basis $z_1^{n_1} z_2^{n_2} z_3^{n_3}$:

$$\hat{\mathcal{L}}(l_{12}^+) z_1^{n_1} z_2^{n_2} z_3^{n_3} = -\lambda q^{2-n_2-n_3+(j_1-j_2)/3} [n_2] z_1^{n_1} z_2^{n_2-1} z_3^{n_3} + \lambda q^{-n_2} [n_3-j_1] z_1^{n_1} z_2^{n_2} z_3^{n_3+1} \quad (4.23)$$

$$\hat{\mathcal{L}}(l_{23}^+) z_1^{n_1} z_2^{n_2} z_3^{n_3} = \lambda q^{n_1+(j_2-j_1)/3} [n_3] z_1^{n_1+1} z_2^{n_2} z_3^{n_3-1} + \lambda [n_1+n_2-n_3-j_2] z_1^{n_1} z_2^{n_2+1} z_3^{n_3} \quad (4.24)$$

$$\begin{aligned} \hat{\mathcal{L}}(l_{13}^+) z_1^{n_1} z_2^{n_2} z_3^{n_3} = & \lambda q^{-n_1-n_2+(2j_2+j_1)/3} [j_1-n_3] z_1^{n_1} z_2^{n_2+1} z_3^{n_3+1} \\ & + \lambda q^{-n_2+2j_2} [n_1-n_2-n_3-j_1+j_2] z_1^{n_1+1} z_2^{n_2} z_3^{n_3} \end{aligned} \quad (4.25)$$

$$\hat{\mathcal{L}}(l_{11}^+) z_1^{n_1} z_2^{n_2} z_3^{n_3} = q^{-n_1-n_3+(2j_1+j_2)/3} z_1^{n_1} z_2^{n_2} z_3^{n_3} \quad (4.26)$$

$$\hat{\mathcal{L}}(l_{22}^+) z_1^{n_1} z_2^{n_2} z_3^{n_3} = q^{-n_2+n_3+(j_2-j_1)/3} z_1^{n_1} z_2^{n_2} z_3^{n_3} \quad (4.27)$$

$$\hat{\mathcal{L}}(l_{33}^+) z_1^{n_1} z_2^{n_2} z_3^{n_3} = q^{n_1+n_2-(j_1+2j_2)/3} z_1^{n_1} z_2^{n_2} z_3^{n_3} \quad (4.28)$$

$$\hat{\mathcal{L}}(l_{21}^-) z_1^{n_1} z_2^{n_2} z_3^{n_3} = \lambda q^{n_1} [n_3] z_1^{n_1} z_2^{n_2} z_3^{n_3-1} + \lambda q^{n_2+n_3+(j_1-j_2)/3} [n_1] z_1^{n_1-1} z_2^{n_2} z_3^{n_3} \quad (4.29)$$

$$\hat{\mathcal{L}}(l_{32}^-) z_1^{n_1} z_2^{n_2} z_3^{n_3} = \lambda [n_2] z_1^{n_1} z_2^{n_2-1} z_3^{n_3} \quad (4.30)$$

$$\hat{\mathcal{L}}(l_{31}^-) z_1^{n_1} z_2^{n_2} z_3^{n_3} = \lambda q^{n_2} [n_1] z_1^{n_1-1} z_2^{n_2} z_3^{n_3} \quad (4.31)$$

These expressions define the representations $\hat{\mathcal{L}}_{\bar{j}}$ which act in the (same for all $\bar{j}$) space $\hat{\mathcal{T}}_{\bar{j}}$ of polynomials $\hat{\phi}$ in z_i (which as a linear space is just T_{00}). (Note however that the concrete form of z_i depends on $\bar{j}$; we do not indicate this dependence explicitly in order to simplify the notation). Using the operations M_i , rescaling the variable $z_i \rightarrow qz_i$, the q-derivative $D_i f = (M_i - M_i^{-1})f/(q - q^{-1})z$ and the operator Z_i , multiplying by the variable z_i , all of which by convention act *directly* on the i^{th} place in the ordered monomials, we obtain the following reduced q-differential representation :

$$\hat{\mathcal{L}}(l_{12}^+) = -\lambda q^{2+(j_1-j_2)/3} M_2^{-1} M_3^{-1} D_2 + \lambda (q^{-j_1} M_3 - q^{j_1} M_3^{-1}) M_2^{-1} Z_3 \quad (4.32)$$

$$\hat{\mathcal{L}}(l_{23}^+) = -\lambda q^{(j_2-j_1)/3} M_1 Z_1 D_3 + \lambda (q^{-j_2} M_1^{-1} M_2^{-1} M_3 - q^{j_2} M_1 M_2 M_3^{-1}) Z_2 \quad (4.33)$$

$$\begin{aligned} \hat{\mathcal{L}}(l_{13}^+) = & \lambda q^{(2j_2+j_1)/3} (q^{j_1} M_3^{-1} - q^{j_1} M_3) M_1^{-1} M_2^{-1} Z_2 Z_3 \\ & + \lambda q^{2j_2} (q^{j_2-j_1} M_1 M_2^{-1} M_3^{-1} - q^{j_1-j_2} M_1^{-1} M_2 M_3) M_2^{-1} Z_1 \end{aligned} \quad (4.34)$$

$$\hat{\mathcal{L}}(l_{11}^+) = q^{(2j_1+j_2)/3} M_1^{-1} M_3^{-1} \quad (4.35)$$

$$\hat{\mathcal{L}}(l_{22}^+) = q^{(j_2-j_1)/3} M_2^{-1} M_3 \quad (4.36)$$

$$\hat{\mathcal{L}}(l_{33}^+) = q^{-(j_1+2j_2)/3} M_1 M_2 \quad (4.37)$$

$$\hat{\mathcal{L}}(l_{21}^-) = \lambda M_1 D_3 + \lambda q^{(j_1-j_2)/3} M_2 M_3^{-1} D_1 \quad (4.38)$$

$$\hat{\mathcal{L}}(l_{32}^-) = \lambda D_2 \quad (4.39)$$

$$\hat{\mathcal{L}}(l_{31}^-) = \lambda M_2 D_1 \quad (4.40)$$

We remark, that although originally we have assumed $j_1, j_2 \in \mathbb{Z}$, the formulae (4.32)-(4.40) define a representation $\hat{\mathcal{T}}_{\bar{j}}$ of $sl_q(3)$ also for arbitrary $j_1, j_2 \in \mathcal{C}$. However, only for $j_1, j_2 \in \mathbb{Z}$ they comprise a deformation of representations which are integrable to a representation of the group $SL(3, \mathcal{C})$.

Now we consider the right regular representation (A 16-18) of the remaining generators on $\phi_j^{\bar{n}}$ and restrict it. It turns out that for any j_1, j_2 there are always unwanted terms like m_{12} or m_{22} in the image of $\mathcal{R}(l_{31}^-)$ and we do not consider this operator any more. Instead, we have

$$\begin{aligned} \mathcal{R}(l_{21}^-) \phi_j^{\bar{n}} = & \lambda q^{(2j_1-j_1)/3} (q^{n_3-1} [n_3] w_1^{n_1} w_2^{n_2} w_3^{n_3-1} m_{11}^{j_1-2} t_{33}^{j_2+1} \\ & - [j_1] w_1^{n_1} w_2^{n_2} w_3^{n_3} m_{11}^{j_1-1} t_{33}^{j_2} m_{12}) , \end{aligned} \quad (4.41)$$

$$\begin{aligned} \mathcal{R}(l_{32}^-) \phi_j^{\bar{n}} = & -\lambda q^{(j_2-j_1)/3} (q^{n_1+n_2-n_3-1} [n_1] w_1^{n_1-1} w_2^{n_2} w_3^{n_3+1} m_{11}^{j_1+1} t_{33}^{j_2-2} \\ & + q^{2n_1+n_2-n_3-1} [n_2] w_1^{n_1} w_2^{n_2-1} w_3^{n_3} m_{11}^{j_1+1} t_{33}^{j_2-2} \\ & + [j_2] w_1^{n_1} w_2^{n_2} w_3^{n_3} m_{11}^{j_1} t_{33}^{j_2-1} t_{32}) . \end{aligned} \quad (4.42)$$

Thus for $\mathcal{R}(l_{21}^-)$ and for $\mathcal{R}(l_{32}^-)$ the unwanted terms (with t_{32} or m_{12}) are not present precisely when $j_1 = 0$ and $j_2 = 0$, respectively. Then, in fact, we obtain two intertwiners: the restriction of $\mathcal{R}(l_{21}^-)$ to $\mathcal{T}_{0,j_2}$ which maps into $\mathcal{T}_{-2,j_2+1}$ and the restriction of $\mathcal{R}(l_{32}^-)$ to $\mathcal{T}_{j_1,0}$ which maps into $\mathcal{T}_{j_1+1,-2}$. By similar computations (first for the squares of these operators and then by induction) one obtains a family of other intertwiners

$$\mathcal{R}(l_{21}^-)^{j_1+1} : \mathcal{T}_{j_1,j_2} \rightarrow \mathcal{T}_{-j_1-2,j_2+j_1+1}, \text{ for } j_1 \in \mathbb{Z}_+, j_2 \in \mathcal{C} \quad (4.43)$$

$$\mathcal{R}(l_{32}^-)^{j_2+1} : \mathcal{T}_{j_1,j_2} \rightarrow \mathcal{T}_{j_1+j_2+1,-j_2-2}, \text{ for } j_2 \in \mathbb{Z}_+, j_1 \in \mathcal{C} \quad (4.44)$$

There are also some mixed intertwiners :

$$\mathcal{R}(l_{32}^-)^{j_2+1} \mathcal{R}(l_{21}^-)^{j_1+1} : \mathcal{T}_{j_1,j_2-j_1-1} \rightarrow \mathcal{T}_{j_2-j_1-1,-j_2-2}, \text{ for } j_1, j_2 \in \mathbb{Z}_+, \quad (4.45)$$

$$\mathcal{R}(l_{21}^-)^{j_1+1} \mathcal{R}(l_{32}^-)^{j_2+1} : \mathcal{T}_{j_1-j_2-1, j_2} \rightarrow \mathcal{T}_{-j_1-2, j_1-j_2-1}, \text{ for } j_1, j_2 \in \mathbb{Z}_+. \quad (4.46)$$

In the same way as for the left-represented operators we perform the explicit reduction to the space $\hat{\mathcal{T}}_{\bar{j}}$ of functions of the variables z_i . Keeping track of the changes of indices $\bar{n}$ and of the labels $\bar{j}$, on which the coefficients C_i depend, we obtain the two basic intertwiners

$$\hat{\mathcal{R}}(l_{21}^-) z_1^{n_1} z_2^{n_2} z_3^{n_3} = \lambda q^{1+2j_2/3} D_3, \quad (4.47)$$

$$\hat{\mathcal{R}}(l_{32}^-) z_1^{n_1} z_2^{n_2} z_3^{n_3} = -\lambda q^{1-j_1/3} D_1 Z_3 - \lambda q^{1-2j_1/3} M_1 D_2. \quad (4.48)$$

The other intertwiners can be easily obtained by observing that the explicit formula for the intertwiner corresponding to product of (any number of) l_{21}^- and l_{32}^- is just the product of the basic intertwiners

$$\hat{\mathcal{R}}((l_{32}^-)^{j_2} (l_{21}^-)^{j_1}) = (\hat{\mathcal{R}}(l_{32}^-))^{j_2} (\hat{\mathcal{R}}(l_{21}^-))^{j_1}. \quad (4.49)$$

They clearly satisfy

$$(\hat{\mathcal{R}}(l_{21}^-))^{j_1+1} \cdot \hat{\mathcal{L}}_{j_1, j_2}(l_{k\ell}^\pm) = \hat{\mathcal{L}}_{-j_1-2, j_2+j_1+1}(l_{k\ell}^\pm) \cdot (\hat{\mathcal{R}}(l_{21}^-))^{j_1+1}, \quad (4.50)$$

$$(\hat{\mathcal{R}}(l_{32}^-))^{j_2+1} \cdot \hat{\mathcal{L}}_{j_1, j_2}(l_{k\ell}^\pm) = \hat{\mathcal{L}}_{j_1+j_2+1, -j_2-2}(l_{k\ell}^\pm) \cdot (\hat{\mathcal{R}}(l_{32}^-))^{j_2+1}, \quad (4.51)$$

$$\hat{\mathcal{R}}((l_{21}^-)^{j_1+1} (l_{32}^-)^{j_2+1}) \cdot \hat{\mathcal{L}}_{j_1-j_2-1, j_2}(l_{k\ell}^\pm) = \hat{\mathcal{L}}_{-j_1-2, j_1-j_2-1}(l_{k\ell}^\pm) \cdot \hat{\mathcal{R}}((l_{21}^-)^{j_1+1} (l_{32}^-)^{j_2+1}), \quad (4.52)$$

$$\hat{\mathcal{R}}((l_{32}^-)^{j_2+1} (l_{21}^-)^{j_1+1}) \cdot \hat{\mathcal{L}}_{j_1, j_2-j_1-1}(l_{k\ell}^\pm) = \hat{\mathcal{L}}_{j_2-j_1-1, -j_2-2}(l_{k\ell}^\pm) \cdot \hat{\mathcal{R}}((l_{32}^-)^{j_2+1} (l_{21}^-)^{j_1+1}), \quad (4.53)$$

where we have explicitly indicated the labels of the representations $\hat{\mathcal{L}}$. These intertwiners give rise to a various partial equivalences between some $\hat{\mathcal{T}}_{\bar{j}}$. In particular, the kernels of these intertwiners form invariant subrepresentations. Thus, for generic $j_1, j_2 \in \mathcal{C}$ the representations $\hat{\mathcal{T}}_{\bar{j}}$ are irreducible except for $j_1 \in \mathbb{Z}_+$ or $j_1 \in \mathbb{Z}_+$, when they are reducible. To study further the reducibility, we note that all the intertwiners which act on $\hat{\mathcal{T}}_{\bar{j}}$ for given $j_1, j_2 \in \mathbb{Z}_+$, contain as a last factor either the $j_1 + 1$ power of the basic intertwiner (4.47) or the $j_2 + 1$ power of (4.48). Thus it is sufficient to consider only the kernels of two intertwiners (4.50) and (4.51). It turns out that their common intersection are irreducible (sub)representations and have the dimension $d = (j_1+1)(j_2+1)(1+(j_1+j_2)/2)$. They consist of polynomials in z_i of limited order, depending on i and on $\bar{j}$, and are q-deformation of the well known representations of $sl(3)$. In particular, $\{1, z_1, z_2\}$ is a basis of $3_q = \ker \hat{\mathcal{R}}(l_{21}^-) \cap \ker(\hat{\mathcal{R}}(l_{32}^-))^2 \subset \hat{\mathcal{T}}_{0,1}$, $\{1, z_3, z_2 z_3 - q^{-1/3} z_1\}$ is a basis of $3_q^* = \ker(\hat{\mathcal{R}}(l_{21}^-))^2 \cap \ker \hat{\mathcal{R}}(l_{32}^-) \subset \hat{\mathcal{T}}_{1,0}$ and $\{1, z_1, z_2, z_3, z_1 z_3, z_2 z_3, q^{1/3}(1+q)z_2^2 z_3 - [2]z_1 z_2, q^{1/3}[2]z_1 z_2 z_3 - (1+q)z_1^2\}$ is a basis of $8_q = \ker(\hat{\mathcal{R}}(l_{21}^-))^2 \cap \ker(\hat{\mathcal{R}}(l_{32}^-))^2 \subset \hat{\mathcal{T}}_{1,1}$.

5. Final remarks.

In this paper, the method previously applied to simpler examples, has been extended to $SL_q(3)$. It permits to obtain in a canonical way some important (reducible and irreducible) representations of $sl_q(3)$ and their intertwiners, and links intimately to what should be a q-analogue of the (infinitesimal) Borel-Weil construction. The reduction procedure corresponds in fact to working on the quantum three dimensional complex flag manifold $F_q(1, 2; 3)$ (the quotient manifold of $SL_q(3)$ by the Borel subgroup), c.f. [5], [6]. This is indeed the case for the representations inside the Weyl chamber ($j_1 \cdot j_2 \neq 0$), while the representations with j_1 or $j_2 = 0$ live, as in the previous simpler examples, over the q-projective spaces (or q-grassmannians). Our functions are local representatives of global sections, like e.g. $m_{11}^{j_1} t_{33}^{j_2}$, of some quantum line bundles over $F_q(1, 2; 3)$. This holds over one particular patch, coordinatized by the variables z_i (or w_i), corresponding to the requirement of invertibility of m_{11} and t_{33} .

We mention that this approach which relies on the infinitesimal covariance and the use of representations of g_q , though closely related, seems to be conceptually simpler than what should be a q-analogue of induced representations (i.e. working with the global covariance under the co-representations of G_q), see for example [7], [8], [9], [10]. A detailed comparison of these two methods and the geometry of the quantum homogeneous space $F_q(1, 2; 3)$ will be discussed in the forthcoming paper.

Acknowledgments

We thank Prof. C. Reina for helpful discussions.

Note added The Heisenberg realization of $U_q sl(n)$ in terms of q-difference operators on the flag manifold is presented in [11]. After submitting this paper, we have been also informed about [12], which has overlap with our results.

Appendix.

Denote $s = \frac{1}{3}(n_{11} + n_{12} + \dots + n_{33})$. The action of the representation $\mathcal{L}$ on $t^{\bar{n}} = t_{11}^{n_{11}} t_{12}^{n_{12}} \dots t_{33}^{n_{33}}$ is given as follows:

$$\begin{aligned} \mathcal{L}(l_{12}^+)t^{\bar{n}} = & -\lambda q^s (q^{1-n_{12}-n_{13}-n_{21}}[n_{21}]t_{11}^{n_{11}+1} \dots t_{21}^{n_{21}-1} \dots t_{33}^{n_{33}} \\ & + q^{1-n_{21}-n_{22}-n_{13}}[n_{22}]t_{11}^{n_{11}} t_{12}^{n_{12}+1} \dots t_{22}^{n_{22}-1} \dots t_{33}^{n_{33}} \\ & + q^{1-n_{21}-n_{22}-n_{23}}[n_{23}]t_{11}^{n_{11}} \dots t_{13}^{n_{13}+1} \dots t_{23}^{n_{23}-1} \dots t_{33}^{n_{33}}) \end{aligned} \quad (A1)$$

$$\begin{aligned} \mathcal{L}(l_{23}^+)t^{\bar{n}} = & -\lambda q^s (q^{1-n_{22}-n_{23}-n_{31}}[n_{21}]t_{11}^{n_{11}} \dots t_{21}^{n_{21}+1} \dots t_{31}^{n_{31}-1} \dots t_{33}^{n_{33}} \\ & + q^{1-n_{23}-n_{31}-n_{32}}[n_{32}]t_{11}^{n_{11}} \dots t_{22}^{n_{22}+1} \dots t_{32}^{n_{32}-1} t_{33}^{n_{33}} \\ & + q^{1-n_{21}-n_{32}-n_{33}}[n_{33}]t_{11}^{n_{11}} \dots t_{23}^{n_{23}+1} \dots t_{33}^{n_{33}-1}) \end{aligned} \quad (A2)$$

$$\begin{aligned} \mathcal{L}(l_{13}^+)t^{\bar{n}} = & -\lambda q^{1-n_{31}+s} (q^{-n_{12}-n_{13}-n_{21}}[n_{31}]t_{11}^{n_{11}+1} \dots t_{31}^{n_{31}-1} \dots t_{33}^{n_{33}} \\ & + q^{-n_{13}-n_{22}-n_{32}}[n_{32}]t_{11}^{n_{11}} t_{12}^{n_{12}+1} \dots t_{32}^{n_{32}-1} t_{33}^{n_{33}} \\ & + q^{-n_{23}-n_{32}-n_{33}}[n_{33}]t_{11}^{n_{11}} \dots t_{13}^{n_{13}+1} \dots t_{33}^{n_{33}-1}) \\ & + \lambda^2 q^{1-n_{31}+s} (q^{1-n_{21}-n_{22}-n_{13}-n_{12}}[n_{31}][n_{22}]t_{11}^{n_{11}+1} \dots t_{21}^{n_{21}+1} t_{22}^{n_{22}-1} \dots t_{31}^{n_{31}-1} \dots t_{33}^{n_{33}} \\ & + q^{1-n_{22}-n_{23}}[n_{31}][n_{23}]t_{11}^{n_{11}} \dots t_{13}^{n_{13}+1} t_{21}^{n_{21}+1} \dots t_{23}^{n_{23}-1} \dots t_{31}^{n_{31}-1} \dots t_{33}^{n_{33}} \\ & + q^{1-n_{23}-n_{32}}[n_{23}][n_{32}]t_{11}^{n_{11}} \dots t_{13}^{n_{13}+1} \dots t_{22}^{n_{22}+1} t_{23}^{n_{23}-1} \dots t_{32}^{n_{32}-1} t_{33}^{n_{33}}) \end{aligned} \quad (A3)$$

$$\mathcal{L}(l_{11}^+)t^{\bar{n}} = q^{s-n_{11}-n_{12}-n_{13}}t^{\bar{n}} \quad (A4)$$

$$\mathcal{L}(l_{22}^+)t^{\bar{n}} = q^{s-n_{21}-n_{22}-n_{23}}t^{\bar{n}} \quad (A5)$$

$$\mathcal{L}(l_{33}^+)t^{\bar{n}} = q^{s-n_{31}-n_{32}-n_{33}}t^{\bar{n}} \quad (A6)$$

$$\begin{aligned} \mathcal{L}(l_{21}^-)t^{\bar{n}} = & \lambda q^{-s} (q^{n_{11}+n_{12}+n_{23}}[n_{13}]t_{11}^{n_{11}} \dots t_{13}^{n_{13}-1} \dots t_{23}^{n_{23}+1} t_{33}^{n_{33}} \\ & + q^{n_{11}+n_{22}+n_{23}}[n_{12}]t_{11}^{n_{11}} t_{12}^{n_{12}-1} \dots t_{22}^{n_{22}+1} \dots t_{33}^{n_{33}} \\ & + q^{n_{21}+n_{22}+n_{23}}[n_{11}]t_{11}^{n_{11}-1} \dots t_{21}^{n_{21}+1} \dots t_{33}^{n_{33}}) \end{aligned} \quad (A7)$$

$$\begin{aligned} \mathcal{L}(l_{32}^-)t^{\bar{n}} = & \lambda q^{-s} (q^{n_{21}+n_{22}+n_{33}}[n_{23}]t_{11}^{n_{11}} \dots t_{23}^{n_{23}-1} \dots t_{33}^{n_{33}+1} \\ & + q^{n_{21}+n_{32}+n_{33}}[n_{22}]t_{11}^{n_{11}} \dots t_{22}^{n_{22}-1} \dots t_{32}^{n_{32}+1} t_{33}^{n_{33}} \\ & + q^{n_{31}+n_{32}+n_{33}}[n_{21}]t_{11}^{n_{11}} \dots t_{21}^{n_{21}-1} \dots t_{31}^{n_{31}+1} \dots t_{33}^{n_{33}}) \end{aligned} \quad (A8)$$

$$\begin{aligned}
\mathcal{L}(l_{31}^-)t^{\bar{n}} &= \lambda q^{n_{33}-s} (q^{n_{11}+n_{12}+n_{23}} [n_{13}] t_{11}^{n_{11}} \dots t_{13}^{n_{13}-1} \dots t_{33}^{n_{33}+1} \\
&\quad + q^{n_{11}+n_{22}+n_{32}} [n_{12}] t_{11}^{n_{11}} t_{12}^{n_{12}-1} \dots t_{32}^{n_{32}+1} t_{33}^{n_{33}} \\
&\quad + q^{n_{21}+n_{31}-n_{32}} [n_{11}] t_{11}^{n_{11}-1} \dots t_{31}^{n_{31}+1} \dots t_{33}^{n_{33}}) \\
&\quad + \lambda^2 q^{n_{33}-s} (q^{n_{11}+n_{22}} [n_{12}] [n_{23}] t_{11}^{n_{11}} \dots t_{12}^{n_{12}-1} \dots t_{22}^{n_{22}+1} t_{23}^{n_{23}-1} \dots t_{33}^{n_{33}+1} \\
&\quad + q^{n_{21}+n_{22}} [n_{11}] [n_{23}] t_{11}^{n_{11}-1} \dots t_{21}^{n_{21}+1} \dots t_{23}^{n_{23}-1} \dots t_{33}^{n_{33}+1} \\
&\quad + q^{n_{21}+n_{32}} [n_{11}] [n_{22}] t_{11}^{n_{11}-1} \dots t_{21}^{n_{23}+1} t_{22}^{n_{22}-1} \dots t_{32}^{n_{32}+1} t_{33}^{n_{33}})
\end{aligned} \tag{A9}$$

$$\begin{aligned}
\mathcal{R}(l_{12}^+)t^{\bar{n}} &= \lambda q^{n_{32}-s} (q^{n_{11}+n_{21}} [n_{31}] t_{11}^{n_{11}} \dots t_{31}^{n_{31}-1} t_{32}^{n_{32}+1} t_{33}^{n_{33}} \\
&\quad + q^{n_{11}+n_{22}} [n_{21}] t_{11}^{n_{11}} \dots t_{21}^{n_{21}-1} t_{22}^{n_{22}+1} \dots t_{33}^{n_{33}} \\
&\quad + q^{n_{12}+n_{22}} [n_{11}] t_{11}^{n_{11}-1} t_{12}^{n_{12}+1} \dots t_{33}^{n_{33}})
\end{aligned} \tag{A10}$$

$$\begin{aligned}
\mathcal{R}(l_{23}^+)t^{\bar{n}} &= \lambda q^{n_{33}-s} (q^{n_{12}+n_{22}} [n_{32}] t_{11}^{n_{11}} \dots t_{32}^{n_{32}-1} t_{33}^{n_{33}+1} \\
&\quad + q^{n_{12}+n_{23}} [n_{22}] t_{11}^{n_{11}} \dots t_{22}^{n_{22}-1} t_{23}^{n_{23}+1} \dots t_{33}^{n_{33}} \\
&\quad + q^{n_{13}+n_{23}} [n_{12}] t_{11}^{n_{11}} t_{12}^{n_{12}-1} t_{13}^{n_{13}+1} \dots t_{33}^{n_{33}})
\end{aligned} \tag{A11}$$

$$\begin{aligned}
\mathcal{R}(l_{13}^+)t^{\bar{n}} &= \lambda q^{n_{33}-s} (q^{n_{11}+n_{21}+n_{32}} [n_{31}] t_{11}^{n_{11}} \dots t_{31}^{n_{31}-1} \dots t_{33}^{n_{33}+1} \\
&\quad + q^{n_{11}+n_{22}+n_{23}} [n_{21}] t_{11}^{n_{11}} \dots t_{21}^{n_{21}-1} \dots t_{23}^{n_{23}+1} \dots t_{33}^{n_{33}} \\
&\quad + q^{n_{13}+n_{12}+n_{23}} [n_{11}] t_{11}^{n_{11}-1} \dots t_{13}^{n_{13}+1} \dots t_{33}^{n_{33}}) \\
&\quad + \lambda^2 q^{n_{33}-s} (q^{n_{11}+n_{22}} [n_{21}] [n_{32}] t_{11}^{n_{11}} \dots t_{21}^{n_{21}-1} t_{22}^{n_{22}+1} \dots t_{32}^{n_{32}-1} \dots t_{33}^{n_{33}+1} \\
&\quad + q^{n_{12}+n_{22}} [n_{11}] [n_{32}] t_{11}^{n_{11}-1} t_{12}^{n_{12}+1} \dots t_{32}^{n_{32}-1} t_{33}^{n_{33}+1} \\
&\quad + q^{n_{12}+n_{23}} [n_{11}] [n_{22}] t_{11}^{n_{11}-1} t_{12}^{n_{12}+1} \dots t_{22}^{n_{22}-1} t_{23}^{n_{23}+1} \dots t_{33}^{n_{33}})
\end{aligned} \tag{A12}$$

$$\mathcal{R}(l_{11}^+)t^{\bar{n}} = q^{-s+n_{11}+n_{21}+n_{31}} t^{\bar{n}} \tag{A13}$$

$$\mathcal{R}(l_{22}^+)t^{\bar{n}} = q^{s+n_{12}+n_{22}+n_{33}} t^{\bar{n}} \tag{A14}$$

$$\mathcal{R}(l_{33}^+)t^{\bar{n}} = q^{s+n_{13}+n_{23}+n_{33}} t^{\bar{n}} \tag{A15}$$

$$\begin{aligned}
\mathcal{R}(l_{21}^-)t^{\bar{n}} &= -\lambda q^{1-n_{12}+s} (q^{-n_{21}-n_{31}} [n_{12}] t_{11}^{n_{11}+1} t_{12}^{n_{12}-1} \dots t_{33}^{n_{33}} \\
&\quad + q^{-n_{22}-n_{31}} [n_{22}] t_{11}^{n_{11}} \dots t_{21}^{n_{21}+1} t_{22}^{n_{22}-1} \dots t_{33}^{n_{33}} \\
&\quad + q^{-n_{22}-n_{32}} [n_{32}] t_{11}^{n_{11}} \dots t_{31}^{n_{31}+1} \dots t_{32}^{n_{32}-1} t_{33}^{n_{33}})
\end{aligned} \tag{A16}$$

$$\begin{aligned}
\mathcal{R}(l_{32}^-)t^{\bar{n}} &= -\lambda q^{1-n_{13}+s} (q^{-n_{22}-n_{32}} [n_{13}] t_{11}^{n_{11}} t_{12}^{n_{12}+1} t_{13}^{n_{13}-1} \dots t_{33}^{n_{33}} \\
&\quad + q^{-n_{23}-n_{32}} [n_{23}] t_{11}^{n_{11}} \dots t_{22}^{n_{22}+1} t_{23}^{n_{23}-1} \dots t_{33}^{n_{33}} \\
&\quad + q^{-n_{23}-n_{33}} [n_{33}] t_{11}^{n_{11}} \dots t_{32}^{n_{32}+1} t_{33}^{n_{33}-1})
\end{aligned} \tag{A17}$$

$$\begin{aligned}
\mathcal{R}(l_{31}^-)t^{\bar{n}} = & -\lambda q^{1-n_{13}+s} \left(q^{-n_{12}-n_{21}-n_{31}} [n_{13}] t_{11}^{n_{11}+1} \dots t_{13}^{n_{13}-1} \dots t_{33}^{n_{33}} \right. \\
& + q^{-n_{22}-n_{23}-n_{31}} [n_{23}] t_{11}^{n_{11}} \dots t_{21}^{n_{21}+1} \dots t_{23}^{n_{23}-1} \dots t_{33}^{n_{33}} \\
& + q^{-n_{23}-n_{32}-n_{33}} [n_{33}] t_{11}^{n_{11}} \dots t_{31}^{n_{31}+1} \dots t_{33}^{n_{33}-1} \left. \right) \\
& + \lambda^2 q^{2-n_{13}+s} \left(q^{-n_{22}-n_{31}} [n_{13}] [n_{22}] t_{11}^{n_{11}} t_{12}^{n_{12}+1} t_{13}^{n_{13}-1} t_{21}^{n_{21}+1} t_{22}^{n_{22}-1} \dots t_{33}^{n_{33}} \right. \\
& + q^{-n_{22}-n_{32}} [n_{13}] [n_{32}] t_{11}^{n_{11}} t_{12}^{n_{12}+1} t_{13}^{n_{13}-1} \dots t_{31}^{n_{31}+1} t_{32}^{n_{32}-1} t_{33}^{n_{33}} \\
& + q^{-n_{23}-n_{32}} [n_{23}] [n_{32}] t_{11}^{n_{11}} \dots t_{22}^{n_{22}+1} t_{23}^{n_{23}-1} t_{31}^{n_{31}+1} t_{32}^{n_{32}-1} t_{33}^{n_{33}} \left. \right) .
\end{aligned} \tag{A18}$$

References

- [1] V.K. Dobrev, Rep. Math. Phys. **25** (1988) 159.
- [2] L. Dąbrowski, V.K. Dobrev, R. Floreanini, J. Math. Phys. **35** (1994) 971.
- [3] L. Dąbrowski, J. Sobczyk, Lett. Math. Phys., in print
- [4] L. D. Faddeev, N. Yu. Reshetikhin and L. A. Takhtajan, Algebraic Analysis **1** (1988) 129.
- [5] Y. Soibelman, RIMS-780, (1991)
- [6] E. Taft, J. Towber, J. Algebra **142** (1991) 1.
- [7] T. Brzeziński, S. Majid, Commun. Math. Phys. **157** (1993) 591.
- [8] R. Budzyński, W. Kondracki, Quantum principal fibre bundles: topological aspects, Warsaw preprint IM PAN 517, 1993.
- [9] M. Durdević, Quantum principal bundles, Belgrade Univ. Preprint, 1993 (hep-th/9311029).
- [10] P. Podleś, Symmetries of quantum spaces. Subgroups and quotient spaces of quantum $SU(2)$ and $SO(3)$ groups, Warsaw Univ. preprint 1994.
- [11] H. Awata, M. Noumi, S. Odake, Lett. Math. Phys. **30** (1994) 35.
- [12] V.K. Dobrev, ASI-TPA/10/93 preprint (hep-th/9405150)